

YUKON SNOW SURVEY BULLETIN & WATER SUPPLY FORECAST

April 1, 2016

Prepared and issued by:
Water Resources Branch
Environment Yukon



PREFACE

The Yukon Snow Survey Bulletin and Water Supply Forecast is prepared and issued three times annually – after April 1st, April 1st and May 1st – by Environment Yukon’s Water Resources Branch. The bulletin provides a summary of winter meteorological and streamflow conditions for Yukon, as well as current snow depth and snow water equivalent observations for 56 locations. This information is used to make projections of total volume runoff for the summer period and an estimate of peak flow for the main river basins and subbasins including the upper and lower Yukon, Pelly, Stewart, Liard, Alsek, Porcupine and Peel Rivers. Information about the bulletin, snowpack conditions or streamflow projections can be obtained by contacting:

David Barrett
Hydrology Technologist
(867) 667-3411
david.barrett@gov.yk.ca

Richard Janowicz
Manager, Hydrology
(867) 667-3223
richard.janowicz@gov.yk.ca

Water Resources Branch, Environment Yukon
(867) 667-3171, toll free (in Yukon): 1-800-661-0408 local 3171
Fax: 867-667-3195 | Email: water.resources@gov.yk.ca

WATER NETWORK CHANGES for 2016

There were no water network changes in 2016. This bulletin, as well as earlier editions, is available online at: www.env.gov.yk.ca/snowbulletin

ISSN 1705-883X

It is recommended that reference to this report be made in the following form:

Yukon Snow Survey Bulletin and Water Supply Forecast, April 1, 2016

© April 2016
Water Resources Branch
Department of Environment
Government of Yukon
Box 2703, Whitehorse, Yukon Y1A 2C6

ACKNOWLEDGMENTS

The Yukon Snow Survey Bulletin and Water Supply Forecast is published three times annually: after April 1st, April 1st, and May 1st. The Bulletin forms part of the Yukon Snow Survey Program administered by the Water Resources Branch, Department of Environment, Government of Yukon.

Other agencies that contribute significantly to the Snow Survey Program by providing data, assistance and information for the bulletin are:

Data Collection Officer, Natural Resources Conservation Service, United States Department of Agriculture

Meteorologist, Wildland Fire Management, Yukon Department of Community Services, Whitehorse

Officer in Charge, Water Survey of Canada, Whitehorse

Water Management Engineer, Yukon Energy Corporation

Agencies cooperating with Environment Yukon in the Snow Survey Program are:

B.C. Ministry of Environment, Water Stewardship Division,

Parks Canada

Yukon Department of Energy Mines and Resources, Compliance Monitoring and Inspections Branch

Yukon Department of Environment, Information Management and Technology Branch

Yukon Department of Highways and Public Works

YUKON TERRITORY SNOWPACK CONDITIONS AND RUNOFF PROJECTION

WEATHER


The warm winter continued throughout the Yukon in March, with the territory experiencing a series of upper ridges culminating in record breaking temperatures during the final week of the month. Old Crow was influenced by several small bursts of Arctic air, but still ended up nearly six degrees warmer than normal. Precipitation was generally below normal in an already dry month with the exception of Dawson and Whitehorse, both of which received moderated snowfall amounts mid-month along with a series of smaller storms.


SNOWPACK


The April 1st Yukon snowpack was either normal or below. Snowpack values ranged from well below normal in southwestern and southeastern Yukon, below normal in central regions and near normal in northern regions. A small pocket of above normal snowpack was observed in the Dawson area

STREAMFLOW

Streamflow conditions throughout Yukon are generally above normal with the exception of the upper Yukon River which is normal. The Liard River was slightly above normal respectively, while the Alsek, Pelly, Stewart, Peel and Porcupine Rivers are well above normal. Streamflow during this period represents winter base flow, which provides an indication of winter groundwater contributions.








YUKON RIVER BASIN

Snowpack conditions in the Yukon River basin range from normal in the northern portion of the basin to well below normal in southwestern regions. Overall conditions for the Yukon River basin are below normal.



UPPER YUKON RIVER SUBBASIN (SOUTHERN LAKES)



Snowpack conditions in the upper Yukon River watershed, are below normal. Values range from 58 percent of normal at Montana Mountain to 102 percent of normal at Log Cabin. The basin wide average has been estimated to be 78 percent of normal.



WHITEHORSE AREA


Snowpack conditions in the Whitehorse area are well below normal for April 1st. Values range from 58 percent of normal at Montana Mountain to 63 percent of normal at Tagish. An area wide average is estimated to be 58 percent of normal.





YUKON RIVER and MARSH LAKE



The mean elevation of Marsh Lake during March was 654.707 m or 0.08 m below normal. Yukon River at Whitehorse mean discharge during March was 90 percent of normal. Given normal summer meteorological conditions, volume runoff and peak flows for the season are each expected to be 80 and 75 percent of normal, respectively.



PELLY RIVER SUBBASIN


Snowpack conditions in the Pelly River watershed are below normal. Values of snow water equivalent range from 65 percent of normal at Hoole River to 89 percent of normal at Twin Creeks. A basin wide average has been estimated to be 79 percent of normal.

Mean March streamflow for the watershed was 183 percent of normal as indicated by the Pelly River below Vangorda Creek. Given normal summer meteorological conditions, volume runoff and peak flows are expected to be 80 percent and 75 percent of normal, respectively.



No data was reported



PELLEY RIVER BELOW VANGORDA CREEK





STEWART RIVER SUBBASIN

Snowpack conditions in the Stewart River watershed are slightly below normal for April 1st. Values of snow water equivalent range from 73 percent of normal at Mayo Airport to 91 percent of normal at Calumet. A basin wide average has been estimated to be 82 percent of normal.



Mean March streamflow for the watershed was 179 percent of normal as indicated by the Stewart River at the Mouth. Given normal summer meteorological conditions, volume runoff and peak flows for the season are each expected to be 80 and 85 percent of normal, respectively.







CENTRAL YUKON RIVER BASIN (CARMACKS AREA)

Snowpack conditions in the Carmacks area are below normal for April 1st. Values of snow water equivalent range from 59 percent of normal at MacIntosh to 81 percent of normal at Williams Creek. An area wide average has been estimated to be 75 percent of normal.



LOWER YUKON RIVER BASIN (DAWSON AREA)



Snowpack conditions in the Dawson area are above normal for April 1st. Values of snow water equivalent range from 80 percent of normal at King Solomon Dome to 140 percent of normal at Midnite Dome. An area wide average has been estimated to be 108 percent of normal.




LIARD RIVER BASIN

Snowpack conditions within the Liard River watershed are below normal. Values of snow water equivalent range from 65 percent of normal at Watson Lake Airport to 89 percent of normal at Tintina Airstrip. A basin wide average has been estimated to be 80 percent of normal.

Mean March streamflow for the Liard River upstream of Upper Liard was 128 percent of normal. Given normal summer meteorological conditions, volume runoff and peak flows for the season are expected to be 75 percent and 80 percent of normal, respectively.





LIARD RIVER AT UPPER CROSSING




ALSEK RIVER BASIN

Snowpack conditions within the Alsek River watershed are well below normal for April 1st. Values of snow water equivalent range from 54 percent of normal at Canyon Lake to 56 percent of normal at Summit. A basin wide average has been estimated to be 55 percent of normal.

Mean monthly streamflow for April as indicated by the Alsek River above Bates River was 156 percent of normal. The Alsek River is primarily a glacial regime type, which is largely dependent on summer temperatures. Given normal summer meteorological conditions, volume runoff and peak flows for the season are expected to be 50 and 55 percent of normal, respectively.





ALSEK RIVER ABOVE BATES RIVER



PEEL RIVER BASIN

Snowpack conditions in the Peel River watershed are below normal with values of snow water equivalent ranging from 67 percent of normal at Ogilvie to 108 percent of normal at Blackstone. A basin wide average has been estimated to be 88 percent of normal.



Mean monthly streamflow for March as indicated by the Peel River above Canyon Creek station was 180 percent of normal. Given normal summer meteorological conditions, volume runoff and peak flows for the season are expected to be 85 and 90 percent of normal, respectively.



PORCUPINE RIVER BASIN

Snowpack conditions in the Porcupine River watershed are below normal with values of snow water equivalent ranging from 82 percent of normal at Eagle River to 125 percent of normal at Old Crow. A basin wide average has been estimated to be 100 percent of normal.

Mean March streamflow for the basin as indicated by the Porcupine River near the International Boundary is 161 percent of normal. There is no flow forecast available for April 1st.



Drainage Basin and Snow Course

For Sample Date: 2016-04-01

Name	Number	Elev (m)	Date of Survey	This Year		Water Content		Yrs of Rec
				Snow Depth (cm)	Water Content (mm)	Last Year (mm)	Average (mm)	
Alsek River Basin								
Canyon Lake	08AA-SC01	1160	2016-03-30	25.3	50	67	94	37
Alder Creek	08AA-SC02	768	2016-03-23	43.5	88 B	110	160	36
Aishihik Lake	08AA-SC03	945	2016-03-30	21.6	33	56	81	22
Haines Junction Farm	08AA-SC04	610	2016-04-01	13	29	73	100	16
Summit	08AB-SC03	1000	2016-04-01	57.2	148	180	260	36
Yukon River Basin								
Tagish	09AA-SC01	1080	2016-03-31	44.3	92	165	148	40
Montana Mountain	09AA-SC02	1020	2016-03-31	37.3	82	154	143	39
Log Cabin (B.C.)	09AA-SC03	884	2016-03-30	96.4	378	340	377	56
Atlin (B.C)	09AA-SC04	730	2016-03-31	23	46	74	121	51
Mt McIntyre B	09AB-SC01B	1097	2016-03-29	43	83	161	158	40
Whitehorse Airport	09AB-SC02	700	2016-03-29	29.4	60	84	103	49
Meadow Creek	09AD-SC01	1235	2016-03-30	84.1	232	314	285	39
Jordan Lake	09AD-SC02	930	2016-03-30	53.9	127	129	140	29
Morley Lake	09AE-SC01	824	2016-03-30	43.2	125	92	153	28
Mount Berdoe	09AH-SC01	1035	2016-03-30	36.3	80	92	110	40
Satasha Lake	09AH-SC03	1106	2016-03-29	33.7	63	65	100	29
Williams Creek	09AH-SC04	914	2016-03-29	43.7	81	93	102	21
Twin Creeks A	09BA-SC02A	900	2016-03-31	67.6	170	183	191	38
Hoole River	09BA-SC03	1036	2016-03-30	40.1	88	164	141	39
Burns Lake	09BA-SC04	1112	2016-03-31	84.4	209	239	225	30
Finlayson Airstrip	09BA-SC05	988	2016-03-30	34.2	74	119	108	29
Fuller Lake	09BB-SC03	1126	2016-03-31	69	167	190	200	29
Russell Lake	09BB-SC04	1060	2016-03-31	82.8	210	240	232	29
Rose Creek	09BC-SC01	1080	2016-03-29	46.1	98	116	112	22
Mount Nansen	09CA-SC01	1021	2016-03-29	31.7	59	60	81	40
MacIntosh	09CA-SC02	1160	2016-03-29	35.4	60	60	100	40
Burwash Airstrip	09CA-SC03	810	2016-03-31	0	0	25	43	39
Beaver Creek	09CB-SC01	655	2016-03-31	29.8	51	57	87	41
Chair Mountain	09CB-SC02	1067	No Surv			98	98	27
Casino Creek	09CD-SC01	1065	2016-03-29	52.1	90	108	128	38
Pelly Farm	09CD-SC03	472	No Surv			88	79	30
Plata Airstrip	09DA-SC01	830	2016-03-31	62.3	148	211	192	38
Arrowhead Lake	09DA-SC02	1120	No Surv			N.S.	196	18
Withers Lake	09DB-SC01	975	2016-03-31	76.2	187	244	233	30
Rackla Lake	09DB-SC02	1040	2016-03-31	78.6	175	170	192	29

Code "E" - Estimate, Code "B" - Survey date is outside of valid sampling range

Drainage Basin and Snow Course

For Sample Date: 2016-04-01

Name	Number	Elev (m)	Date of Survey	This Year		Water Content		
				Snow Depth (cm)	Water Content (mm)	Last Year (mm)	Average (mm)	Yrs of Rec
Yukon River Basin								
Mayo Airport A	09DC-SC01A	540	2016-03-29	27.8	70	103	97	46
Mayo Airport B	09DC-SC01B	540	2016-03-29	28	48	118	107	28
Edwards Lake	09DC-SC02	830	2016-03-31	58.2	150	156	164	29
Calumet	09DD-SC01	1310	2016-03-29	85.9	181	232	197	37
King Solomon Dome	09EA-SC01	1080	2016-03-30	75.1	128	189	163	41
Grizzly Creek	09EA-SC02	975	2016-03-31	69	167	233	180	40
Midnight Dome	09EB-SC01	855	2016-03-29	83.2	205	181	152	41
Boundary (Alaska)	09EC-SC02	1005	2016-03-30	76.2	165.1	117	135	46
Porcupine River Basin								
Riff's Ridge	09FA-SC01	650	2016-03-31	78.8	136	213	148	28
Eagle Plains	09FB-SC01	710	2016-03-31	86.5	158	223	164	32
Eagle River	09FB-SC02	340	2016-03-31	67.4	113	181	137	32
Old Crow	09FD-SC01	299	2016-03-30	78.6	144	191	119	34
Liard River Basin								
Watson Lake Airport	10AA-SC01	685	2016-03-29	50.9	89	120	141	51
Tintina Airstrip	10AA-SC02	1067	2016-03-30	73.7	184	188	208	38
Pine Lake Airstrip	10AA-SC03	995	2016-03-31	63.8	196	226	229	40
Ford Lake	10AA-SC04	1110	2016-03-30	68.5	149	188	196	29
Frances River	10AB-SC01	730	2016-03-30	52.4	114	113	163	41
Hyland River	10AD-SC01	855	2016-03-30	58.7	137	157	180	39
Peel River Basin								
Blackstone River	10MA-SC01	920	2016-03-31	46.6	115	122	105	40
Ogilvie River	10MA-SC02	595	2016-03-31	66.25	71	139	107	39
Bonnet Plume Lake	10MB-SC01	1120	2016-03-31	71.5	156	154	180	29
Alaska Snow Courses								
Eaglecrest	08AK-SC01	305	2016-03-31	0	0	5	517	34
Moore Creek Bridge	08AK-SC02	700	2016-03-31	106.68	411.48	386	535	24

Code "E" - Estimate, Code "B" - Survey date is outside of valid sampling range


INDEX OF YUKON SNOW COURSES

NAME	NUMBER	ELEVATION (m)	LATITUDE	LONGITUDE	AGENCY
YUKON RIVER BASIN					
Tagish	09AA-SC1	1080	60°17'	134°11'	2
Montana Mountain	09AA-SC2	1020	60°08'	134°44'	2
Log Cabin (B.C.)	09AA-SC3	884	59°46'	134°58'	2
Atlin (B.C.)	09AA-SC4	730	59°34'	133°42'	3
Mt. McIntyre (B)	09AB-SC1B	1097	60°39'	135°08'	1
Whitehorse Airport	09AB-SC2	700	60°42'	135°04'	1
Meadow Creek	09AD-SC1	1235	60°35'	133°05'	2
Jordan Lake	09AD-SC2	930	60°52'	132°50'	2
Morley Lake	09AE-SC1	824	60°00'	132°07'	2
Mount Berdoe	09AH-SC1	1035	62°02'	136°14'	2
Satasha Lake	09AH-SC3	1106	61°29'	136°16'	2
Williams Creek	09AH-SC4	914	60°21'	136°43'	2
Twin Creeks	09BA-SC2	900	62°37'	131°16'	2
Hoole River	09BA-SC3	1036	61°32'	131°36'	2
Burns Lake	09BA-SC4	1112	62°17'	129°57'	2
Finlayson Airstrip	09BA-SC5	988	61°42'	130°46'	2
Fuller Lake	09BB-SC3	1126	62°58'	130°46'	2
Rose Creek	09BC-SC01	1080	62°20'	133°23'	2
Russell Lake	09BB-SC4	1060	63°12'	133°29'	2
Mount Nansen	09CA-SC1	1021	62°02'	137°03'	2
Macintosh	09CA-SC2	1160	61°43'	137°20'	2
Burwash Airstrip	09CA-SC3	810	61°23'	139°03'	2
Duke River	09CA-SC5	1310	61°15'	138°59'	6
Beaver Creek	09CB-SC1	655	62°25'	140°51'	2
Chair Mountain	09CB-SC2	1067	62°04'	140°48'	2
White River	09CB-SC3	823	61°55'	140°32'	2
Casino Creek	09CD-SC1	1164	62°44'	138°48'	2
Pelly Farm	09CD-SC3	472	62°50'	137°20'	8
Plata Airstrip	09DA-SC1	830	63°31'	132°03'	2
Arrowhead Lake	09DA-SC2	1120	63°42'	131°10'	2
Withers Lake	09DB-SC1	975	63°59'	132°18'	2
Rackla Lake	09DB-SC2	1040	64°17'	133°15'	2
Mayo Airport (A)	09DC-SC1A	540	63°38'	135°53'	2
Mayo Airport (B)	09DC-SC1B	540	63°38'	135°53'	2
Edwards Lake	09DC-SC2	830	63°42'	134°18'	2
Calumet	09DD-SC1	1310	63°55'	135°24'	2
King Solomon Dome	09EA-SC1	1080	63°52'	138°56'	2
Grizzly Creek	09EA-SC2	975	64°26'	138°16'	2
Boundary (Alaska)	09EC-SC2	1005	64°05'	141°27'	4
Midnight Dome	09EB-SC1	855	64°04'	139°24'	2

NAME	NUMBER	ELEVATION (m)	LATITUDE	LONGITUDE	AGENCY
LIARD RIVER BASIN					
Watson Lake Airport	10AA-SC1	685	60°07'	128°50'	2
Tintina Airstrip	10AA-SC2	1067	61°05'	131°15'	2
Pine Lake Airstrip	10AA-SC3	995	60°06'	130°56'	2
Ford Lake	10AA-SC4	1110	60°47'	131°28'	2
Frances River	10AB-SC1	730	60°35'	129°11'	2
Hyland River	10AD-SC1	855	61°31'	128°16'	2
ALSEK RIVER BASIN					
Canyon Lake	08AA-SC1	1160	61°07'	136°59'	7
Alder Creek	08AA-SC2	768	60°22'	137°06'	2
Aishihik Lake	08AA-SC3	945	61°12'	137°00'	7
Haines Junction Farm	08AA-SC4	610	60°45'	137°34'	2
Clay Creek	08AB-SC2	670	60°09'	137°56'	6
Summit	08AB-SC3	1000	60°51'	137°47'	2
Profile Mountain	08AB-SC4	900	60°38'	137°56'	6
PEEL RIVER BASIN					
Blackstone River	10MA-SC1	920	64°57'	138°15'	2
Ogilvie River	10MA-SC2	595	65°21'	138°18'	2
Bonnet Plume Lake	10MB-SC1	1120	64°18'	132°00'	2
PORCUPINE RIVER BASIN					
Riff's Ridge	09FA-SC1	650	65°57'	137°22'	2
Eagle Plains	09FB-SC1	710	66°22'	136°44'	2
Eagle River	09FB-SC2	340	66°27'	136°43'	2
Old Crow	09FD-SC1	299	67°34'	139°51'	6
ALASKA SNOW COURSES					
Eaglecrest	34J03	305	58°17'	134°32'	4
Moore Creek Bridge	34K02	701	59°31'	135°15'	4

Numbers refer to Agencies cooperating in the Yukon Snow Surveys:

1. Department of Environment, Government of Yukon
2. Dept of Energy Mines and Resources Yukon
3. British Columbia Ministry of Environment
4. USDA Natural Resources Conservation Service
5. Yukon Transportation and Highways
6. Parks Canada
7. Yukon Energy Corp.
8. Private Contract



Location of Water Resource Snow Courses