

Biweekly respiratory surveillance report for week ending October 21, 2023

- For respiratory surveillance in Canada, visit: [Respiratory Virus Detection Surveillance System and COVID-19 Data Trends](#)
- For surveillance of COVID-19 variants of concern, visit World Health Organization: [Tracking SARS-CoV-2 Variants](#)
- For COVID-19 vaccination coverage in Canada, visit: [COVID-19 vaccination in Canada](#)

Overall Summary

Respiratory virus activity has remained stable in recent weeks in the Yukon. The table below describes the intensity, geographical spread, and local trends.

	Level	Rationale
Intensity	Medium	Activity occurring around historical levels in most indicators, but higher than historical indicators in others (e.g., some regions and Whitehorse COVID-19 hospitalizations)
Geographical spread	Regional	Activity mainly concentrated in Whitehorse area, with some rural areas experiencing higher activity than others
Trend	Stable	Activity is relatively unchanged from last week

The intensity is based on the overall level of clinical activity in the Yukon, measured through syndromic, sentinel, active and lab surveillance indicators. Low means no activity or activity below baseline, medium means expected or average levels of activity, high means levels of activity that are higher than historical baseline, and very high means exceptionally high levels of activity*.

*Baseline is the level at which activity remains throughout summer and most of the winter, or based on historical data for the same time period in previous years, where available.

The geographical spread is rated as either no activity (no evidence of increased or unusual respiratory disease activity), regional (activity occurring in some but not all regions of Yukon), or widespread (activity generally occurring across all of the Yukon).

The trend is a comparison of surveillance indicators to the previous two weeks. Increasing means there is evidence that the level of respiratory disease activity is increasing, stable means respiratory disease activity is relatively unchanged, and decreasing means there is evidence that the level of respiratory disease activity is decreasing.

Disease summaries

The table below provides general trends from the different surveillance indicators for each disease that is included in this report.

	COVID-19	Influenza	RSV
Case rates	Stable	Stable	Stable
Percent positivity	Stable	Stable	Stable
Syndromic indicators		Stable	
Severity	Stable to increasing	Stable	Stable
Wastewater	Stable	N/A	N/A
Vaccination uptake	Stable	Increasing	N/A

Laboratory indicators

- COVID-19 test positivity remained the same compared to the previous two weeks, and was highest in the Whitehorse area
 - Influenza A test positivity remained the same compared to the previous two weeks, and was equal between rural Yukon and the Whitehorse area
 - Influenza B test positivity remained the same compared to the previous two weeks, and was equal between rural Yukon and the Whitehorse area
 - RSV test positivity remained the same compared to the previous two weeks, and equal between rural Yukon and the Whitehorse area
 - COVID-19 was the organism with the highest average test positivity over the last two weeks
 - Testing volume was highest among Whitehorse residents over the last two weeks
 - COVID-19 wastewater viral load in Haines Junction is stable
-

Figure 1: Percent positivity by respiratory virus




Figure 2: Number of COVID-19 tests and percent positivity by pathogen and region




Figure 3: Number of Influenza A tests and percent positivity by region




Figure 4: Number of Influenza B tests and percent positivity by region




Figure 5: Number of RSV tests and percent positivity by region






Figure 6: COVID-19 wastewater surveillance – Haines Junction



Syndromic surveillance indicators

- Visits to community health centres for respiratory-related symptoms were similar for the Central and West regions, and higher for the North and Southeast regions compared to previous years. Respiratory visits were highest in the Southeast region.
- Visits to community health centres for respiratory-related symptoms are increasing in the North and Southeast regions and stable in the Central and West regions.


Figure 7: Percentage of weekly Community Health Centre* visits for acute respiratory-related reasons, by region



Sentinel surveillance indicators

- The rate of calls for respiratory-related reasons to 811 stayed the same compared to the previous two weeks
-

Figure 8: Rate of 811 calls that are respiratory-related



Active surveillance indicators

- The rate of confirmed cases of COVID-19 in the Yukon is stable, and similar to previous years (aside from 2021-2022). Hospitalization rates are stable, and higher than previous years.
- The rate of confirmed cases of Influenza A in the Yukon is stable, and similar to previous years. Hospitalization rates are stable, and similar to previous years.
- The rate of confirmed cases of Influenza B in the Yukon is stable, and similar to previous years. Hospitalization rates are stable, and similar to previous years.

- The rate of confirmed cases of RSV in Yukon is stable, and similar to previous years. Hospitalization rates are stable, and similar to previous years.
- For COVID-19, influenza and RSV combined, case rates in rural Yukon are stable, and similar to previous years. Hospitalization rates are stable, and similar to previous years, except in comparison to 2021-22 during which rates were higher.
- For COVID-19, influenza and RSV combined, case rates in the Whitehorse area are stable, and similar to previous years. Hospitalization rates are increasing, and higher than previous years.

Figure 9: Case rates per 100,000 by respiratory virus




Figure 10: Hospitalization rates per 100,000 by respiratory virus




Table 1: Respiratory illness death rates per 100,000 by virus

Respiratory season	COVID-19 rate	Influenza A rate	Influenza B rate	RSV rate
2020-21	19.0	0	0	0.0
2021-22	51.0	0	0	0.0
2022-23	13.7	0	0	2.3
2023-24	6.7	0	0	0.0



Figure 11: Respiratory illness case rates per 100,000 by residence




Figure 12: Respiratory illness hospitalization rates per 100,000 by residence




Figure 13: Respiratory virus case and hospitalization rates per 100,000, current season





Figure 14: Respiratory virus outbreaks, current season



Immunization Indicators

Book your COVID-19 and flu vaccine appointment here

Immunization indicators are updated monthly in the first full week of the month.

- Influenza vaccination coverage is below 1% for the Yukon population this season.
- Overall, 19% of the Yukon population has had a bivalent COVID-19 vaccination as their most recent dose. The highest bivalent booster coverage is in the 70+ age group.
- Influenza vaccination uptake is increasing.

Figure 15: Influenza vaccine uptake (%), by year




Figure 16: Influenza vaccine uptake (%), current season, by age group




Figure 17: Influenza vaccine uptake (%), current season, by client health region




Figure 18: COVID-19 vaccination status, by age group




Figure 19: Percent of population whose most recent dose was a COVID-19 bivalent vaccination, by client health region and age group





Figure 20: Cumulative uptake of influenza vaccine, by season



Data Notes

All information is subject to change as reconciliation occurs and data becomes more complete.

Epidemiological weeks are standardized ways to count events on weekly basis year after year. The epidemiological weeks used in this report run from Sunday to Saturday. A full calendar can be found on the [Public Health Agency of Canada FluWatch website](#)

This report is updated at the following frequency during respiratory season (late August to early June): laboratory, syndromic, sentinel, and active surveillance indicators updated bi-weekly; immunization indicators updated monthly.

This report is updated at the following frequency outside of respiratory season (early June to late August): laboratory, syndromic, sentinel, and active surveillance indicators updated monthly; immunization indicators not updated.

Lab Indicators

- Based on lab-based PCR tests and rapid point-of-care tests from Whitehorse General Hospital. This does not include at-home rapid tests or tests completed in private labs.
- Percent positivity is based on the volume of specimens. As such, one individual can appear in the data more than once.
- Data will be impacted by clinical testing criteria, availability of testing, and healthcare seeking behaviour.
- Indicators stratified by rural and Whitehorse area residence reflect the information on where the person who was tested resides, and does not necessarily indicate where that person was tested or was located at the time of infection.
- Wastewater surveillance data is extracted from the [Public Health Agency of Canada COVID-19 wastewater surveillance dashboard](#)

Syndromic Surveillance Indicators

- Visits are classified by syndromes, which include clinician assessment and groupings of clinical signs and symptoms. As such, these are not considered diagnoses of communicable disease, and visits that are not related to respiratory viral illness may be included.

- Data is presented at the visit level, as such one individual can appear in the data more than once if they have several visits.
 - Data may be impacted by factors such as healthcare seeking behaviour, and availability of health services.
 - Community health centre visits reflect the regional location of the health centre, and not necessarily the residence of people visiting the health centre.
-

Sentinel Surveillance Indicators

- Includes calls to 811 from Yukon residents for respiratory-related reasons.
 - Data may be impacted by factors such as the public's relative concern for symptoms or illness, and availability of health services.
-

Active Surveillance Indicators

- Includes all Yukon resident cases reported to Yukon Communicable Disease Control. Does not include non-resident cases that were diagnosed in the Yukon.
 - Data will be impacted by clinical testing criteria and availability of testing.
 - Episode dates are based on either date of symptom onset, laboratory collection date, or date of report to public health, depending on availability of data.
 - Geographical region (rural, Whitehorse, etc.) is based on patient residence, not service delivery location.
 - Death rates are estimated from Yukon Communicable Disease Control investigations, and do not reflect finalized Vital Statistics Database data. As such, these rates may be subject to some variation.
 - Outbreaks are reported for institutional settings, including but not limited to hospitals, long term care, correctional facilities, and other congregate settings.
-

Immunization Indicators

- Vaccine status of Health Canada approved COVID-19 vaccines:
- 

- At least one dose
 - Primary series completed
 - Received a bivalent vaccine as their most recent booster dose
 - Received an XBB vaccine as their most recent booster dose
- Client health region/branch indicates the most recently available information on where a person resides and does not necessarily indicate where that person received the vaccine